Nonuniversal critical dynamics on planar random lattices with heterogeneous degree distributions

Sidiney G. Alves,^{1,*} Silvio C. Ferreira,^{2,3,†} and Marcelo M. de Oliveira^{1,‡}

¹Departamento de Estatística, Física e Matemática,

Universidade Federal de São João Del-Rei 36490-972, Ouro Branco, MG, Brazil

²Departamento de Física, Universidade Federal de Viçosa, 36570-900 Viçosa, Minas Gerais, Brazil

³National Institute of Science and Technology for Complex Systems, 22290-180, Rio de Janeiro, Brazil

The weighted planar stochastic (WPS) lattice introduces a topological disorder that emerges from a multifractal structure. Its dual network has a power-law degree distribution and is embedded in a two-dimensional space, forming a planar network. We modify the original recipe to construct WPS networks with degree distributions interpolating smoothly between the original power-law tail, $P(q) \sim q^{-\alpha}$ with exponent $\alpha \approx 5.6$, and a square lattice. We analyze the role of disorder in the modified WPS model, considering the critical behavior of the contact process. We report a critical scaling depending on the network degree distribution. The scaling exponents differ from the standard mean-field behavior reported for CP on infinite-dimensional (random) graphs with powerlaw degree distribution. Furthermore, the disorder present in the WPS lattice model is in agreement with the Luck-Harris criterion for the relevance of disorder in critical dynamics. However, despite the same wandering exponent $\omega = 1/2$, the disorder effects observed for the WPS lattice are weaker than those found for uncorrelated disorder.

I. INTRODUCTION

Nonequilibrium phase transitions between an active (fluctuating) state and an inactive, absorbing state [1, 2] arise frequently in a broad class of phenomena, such as epidemic spreading [3], chemical reactions [4, 5], population dynamics [6], and related fields. Experimental realizations in liquid crystal electroconvection [7], driven suspensions [8], and superconducting vortices [9] also highlight the importance of this kind of transition.

In analogy with equilibrium phase transitions, it is expected that a continuous absorbing phase transition (APT) can be classified into universality classes depending on a few characteristics of the model, namely symmetries, conserved quantities, and dimensionality [2]. Generically, models such as the contact process (CP), which exhibits a continuous APT, belong to the directed percolation (DP) universality class in the absence of extra symmetries, disorder, or conservation laws [2, 10, 11]. Other robust classes emerge when conservation laws are included [2].

Of particular interest is how spatially quenched disorder affects the critical behavior of an APT. Quenched disorder was introduced on regular lattices by random deletion of sites [12-17] or by random spatial fluctuations of the control parameter [18-20]. In these cases, it was shown that quenched randomness produces rare regions that are locally supercritical, even when the whole system is subcritical [21]. The lifetime of such active rare regions grows exponentially with the domain size, leading to slow dynamics with nonuniversal exponents for some interval of the control parameter $\lambda_{\rm c}(0) < \lambda < \lambda_{\rm c}$, where $\lambda_{\rm c}(0)$ and $\lambda_{\rm c}$ are the critical points of the clean (ordered) and disordered systems, respectively. This extended interval where critical behavior is observed is called Griffiths phase (GP), and it was verified in DP models with uncorrelated disorder, irrespective of the disorder strength [13, 14, 16, 17, 20]. Another manifestation of relevant disorder is the smearing of a critical phase transition [15, 22] where the singular critical behavior given by power-law is replaced by stretched exponential with characteristic scales.

The Harris' criterion [23] states that uncorrelated quenched disorder is a relevant perturbation if $d\nu_{\perp} < 2$, where d is the dimensionality and ν_{\perp} is the correlation length exponent of the clean model. In the DP class, where $\nu_{\perp} = 1.096854(4)$, 0.7333(75), and 0.584(5), for d = 1, 2, and 3, respectively [2], the criterion is satisfied for all dimensions d < 4, and, consequently, the disorder is predicted to be relevant according to the Harris criterion. Previous works, based on renormalization group methods and numerical simulations, concluded that the activated-disorder behavior belongs to the universality class of the random transverse Ising model [24, 25], thus confirming the Harris' criterion predictions.

A distinct kind of disorder is introduced by random lattices generated by the Voronoi-Delaunay (VD) triangulation [26], which is a two-dimensional planar graph with a Poissonian distribution of connectivity with an average degree $\bar{q} = 6$. In this case, the disorder does not change the critical behavior exhibited by the clean CP [27, 28], which was, in principle, at odds with the above-mentioned results for the uncorrelated disorder, which led to strong GPs. Barghathi and Vojta [29] proposed a solution to the puzzle by applying the theory of Luck [30]. The Luck-Harris criterion considers that the stability of the critical point is governed by the decay of spatial fluctuations in the local coordination num-

^{*} sidiney@ufsj.edu.br

[†] silviojr@ufv.br

[‡] mmdeoliveira@ufsj.edu.br

bers. Averaging the coordination number q over coarsegrained blocks, the corresponding variance σ_q decays algebraically with the block size $L_{\rm b}$, $\sigma_q \sim \dot{L}_{\rm b}^{-a}$, where $a = d(1 - \omega)$ and ω is the wandering exponent [30]. So, the criterion states that quenched disorder is an irrelevant perturbation if the inequality $a\nu_{\perp} > 1$ holds. In the case of random dilution, the central limit theorem leads to a = d/2, and the criterion reduces to the original Harris criterion. Barghathi and Voita [29] showed that coordination spatial fluctuations in VD lattices and other disordered planar graphs are not governed by uncorrelated disorder (a = 1 in d = 2), but by strong anticorrelations with exponent a = 3/2 such that VD lattices satisfy the Luck-Harris condition for irrelevance of disorder since $a_c = 1/\nu_{\perp} = 1.3636 < 3/2$. However, Schrauth and collaborators [31-33], reported examples that violate the Luck-Harris criterion even using the approach of Barghathi and Vojta [29].

A structure presenting topological disorder was proposed by Hassan et al., the weighted planar stochastic (WPS) lattice [34]. In the WPS lattice, the topological disorder emerges from a multifractal structure [35], with its coordination number probability distribution following a power law. It was found that the isotropic percolation universality class on the WPS lattice [36] is distinct from the one for all the regular planar lattices [37]. An opinion-dynamic model also exhibits a critical behavior on the WPS different from the one observed in regular lattices [38]. Recently, it was shown that the critical behavior of the contact process on the WPS lattice [39] is different from that on a regular one, showing that the coordination disorder introduced by this model is a relevant perturbation of the model.

On the other hand, in infinite-dimensional power-law networks, it is known that the critical exponents depend on the degree distribution exponent [40, 41] consistently: the exponents agree with the heterogeneous mean-field theory, which explicitly takes into account the shape of the degree distribution [40, 41]. In particular, the critical exponents are equivalent to those of the usual mean-field theory for non-scale-free networks with degree exponent $\alpha > 3$, in contrast with the WPS result, where unusual mean-field exponents were found, despite the degree exponent $\alpha \approx 5.6$ out of the scale-free regime.

So, to further investigate the effects of topological planar disorder, we study the critical behavior of the CP in a modified structure based on the WPS lattice. We change the original algorithm of the partitioning process to introduce dual networks with distinct degree distribution exponents $\alpha \geq \alpha_{WPS}$. We analyze the critical behavior of the CP on these structures, which exhibits critical scaling that depends on the lattice degree exponent. This dependence differs from infinite-dimensional powerlaw networks and indicates a generalized DP universality class whose exponents depend on both embedding dimension and degree distribution. Finally, the wandering exponent analysis predicts that the introduced disorder is relevant due to nonuniversal critical exponents that differ from the DP universality class. However, the disorder effects are weaker than those usually observed in uncorrelated cases, where extended regions of criticality are observed [13, 16, 17, 21].

The remainder of this paper is organized as follows. In Sec. II, we present the model and methods employed in the work. In Sec. III, we show and discuss our simulation results. Section IV is to summarize our conclusions.

II. MODELS AND METHODS

A. Modified WPSL

We start by presenting the standard algorithm for the WPS lattice introduced in Ref. [34]. Consider a square domain. In the first step, it is divided randomly into four rectangular blocks. In the following steps, a block is chosen at random, one at a time, with probability proportional to its area, and partitioned into four new blocks at random. After n iterations, the lattice has 3n + 1 nodes. Finally, the dual lattice is constructed in the following way: each block is considered a node of the network, and two blocks are connected if they share borders.

The modified WPS algorithm limits the partitioning region around the squares' centers within some interval controlled by a parameter $\epsilon \in [0, 1]$. At each iteration, a rectangle of sides (ℓ_x, ℓ_y) is chosen proportionally to its area. The partitioning points (x, y) are chosen at random within the domain $(1 - \epsilon)\ell_x/2 < x < (1 + \epsilon)\ell_x/2$ and $(1 - \epsilon)\ell_y/2 < y < (1 + \epsilon)\ell_y/2$, where ϵ is the parameter that controls the randomness. Notice that the limits $\epsilon = 0$ and $\epsilon = 1$ correspond to a square lattice and the original WPS lattice, respectively. By construction, the average area of the blocks is $\langle A \rangle = 1/(3n + 1)$, which defines a typical average length $\langle \ell \rangle = 1/\sqrt{3n + 1}$.

Figure 1(a) shows a snapshot of a random WPS lattice generated after n = 85 iterations (256 nodes). The WPS lattice presents disorder due to its block's multifractal structure, which leads to a coordination number with a distribution that follows a power law, $P(q) \sim q^{-\alpha}$, with $\alpha = -5.6$ [34]. Figures 1(b)-1(d) illustrate different block patterns obtained using three values of $\epsilon = 0.1, 0.5$, and 0.9 after n = 85 iterations of the rule. The bottom panels of the figure 1 show the corresponding dual networks obtained from the block diagrams of the top panels.

The degree distributions obtained with the modified algorithm are presented in Fig. 2. The networks have degree exponents varying with ϵ , which interpolate between the original WPS model with exponent $\alpha \approx 5.6$ and a regular distribution of a square lattice. For $\epsilon \ll 1$, the distributions become Poissonian and lose their power-law tail. Indeed, large decay exponents $\alpha \gg 1$ are consistent with an exponential tail.

We also investigate how the fluctuation of the average connectivity scales with the parameter ϵ using the wandering exponent obtained in the following way: The system is divided into blocks of size $L_{\rm b}$, and the block-



FIG. 1. Illustration of block diagrams (top panels) and the dual network (bottom panels) obtained using the (a) original WPS ($\epsilon = 1$) and the modified one with (b) $\epsilon = 0.1$, (c) 0.5, and (d) 0.9 after n = 85 iterations.

averaged standard deviation σ_q in relation to the overall average degree $\langle q \rangle$ is calculated [29]. Figure 3(a) shows σ_q as a function of the grid size $L_{\rm b}/\langle \ell \rangle$. The exponent $a = d(1-\omega)$ of the scaling law $\sigma_q \sim L_{\rm b}^{-a}$ presents an asymptotic decay for $L_b/\langle \ell \rangle \to \infty$ consistent with uncorrelated disorder for which a = 1. The corresponding wandering exponent satisfies the Luck-Harris criterion for models in the DP class, where disorder is a relevant perturbation of the dynamics, when $a \ll a_c = 1.363$. A curious behavior is present in the initial decay, where an exponent close to the critical value $a_c = 1.363$ is reported. To the best of our understanding, this is a coincidence. Finally, we also analyzed how the average degree with blocks of size $L_{\rm b}$, $\langle q \rangle_b$, approaches the global $\langle q \rangle$ computed for the whole lattice, presented in figure 3(b). A decay given by $\langle q \rangle_b - \langle q \rangle \sim L_{\rm b}^{-3}$ was observed for all values of $\epsilon \in [0.1, 0.9]$. Since this decay is much faster than that of the degree fluctuations, the pitfall indicated in Ref. [29], where the choice of $\langle q \rangle_b$ or $\langle q \rangle$ can alter the wandering exponent, does not apply to our current analysis.

B. Contact process

The CP is a stochastic interacting particle system defined on the top of a lattice [1, 23]. So, each lattice node



FIG. 2. Degree probability distribution for the dual WPS lattices obtained for various values of ϵ . The dashed lines is a power law with exponent equals to -5.63.

i can be in one of the two states: occupied by a particle $(\sigma_i(t) = 1)$ or empty $(\sigma_i(t) = 0)$. The stochastic process evolves via a catalytic creation of particles: a new particle is created at a node *i* with rate λr , where *r* is the fraction of occupied nearest-neighbors of *i*. The CP dynamics also include particle spontaneous annihilation with a rate $\mu = 1$, independently of the neighboring sites. If all nodes are empty, the system is trapped in the ab-



FIG. 3. (a) Fluctuation of the mean degree (or connectivity) as a function of the grid size $L_{\rm b}/\langle \ell \rangle$. The solid and dashed lines represent power law decays with exponent $a = a_c = 1.363$ and a = 1, respectively. (b) Difference between average degree computed for blocks of size $L_{\rm b}$ and the full network.

sorbing state. Exactly at $\lambda = \lambda_c$, the system undergoes a continuous phase transition from the active phase to an absorbing phase within the directed percolation universality class.

The CP simulations on graphs can be implemented as follows [1] (See also [42]): Randomly, select an occupied site. Then, choose between (i) annihilation, with probability $1/(1 + \lambda)$, or (ii) creation, with probability $\lambda/(1+\lambda)$. If annihilation is selected, the node is vacated. On the other hand, if creation is chosen, one of the qnearest-neighbor nodes of the selected node is chosen at random, and if empty, it becomes occupied. Otherwise, no change of state happens. The time increment associated with each event of creation or annihilation is given by $\Delta t = -\ln \xi/[(\mu + \lambda)N_p]$, where N_p is the number of occupied nodes just before the attempted transition and ξ is a random number uniformly distributed in the interval (0, 1).

III. RESULTS AND DISCUSSION

We begin our study by analyzing the evolution of the number of active nodes n(t). This spreading analysis is done by considering the initial condition with a single particle. The critical value of the control parameter λ_c is defined as the smallest λ supporting asymptotic growth of n(t) [1]. We found the critical value depends slightly on the parameter ϵ ; see Table I. In this part, we used lattices with 10⁶ iterations and took an average over 5×10^3 lattices, with 10 runs in each one.

In Fig. 4, we show the time evolution of the number of particles n(t) at $\lambda = \lambda_c$ for different values of ϵ . We observe that the asymptotic evolution follows a power law, $n(t) \propto t^{\eta}$, as expected for critical spreading [1, 2]. The exponent η varies continuously with ϵ , as shown in Table I. The critical exponents for the smallest values of ϵ are closer to the exponent $\eta = 0.2295(10)$ of the DP class [2]. Conversely, the value $\eta = 0.185(5)$ for the original WPS lattice [39] is recovered when $\epsilon \to 1$.



FIG. 4. Number of active sites as a function of time in the spreading analysis. We consider different values of the control parameter. The lines are power-law regressions (Pearson regression coefficient r > 0.999) of the data represented by symbols.

Figure 5 presents de decays of the order parameter for simulations with a fully active initial condition $(\rho(0) = 1)$. At criticality, one expects that the density of active sites $\rho(t)$ decays as [1] $\rho \propto t^{-\delta}$. We found continuously varying values of the critical exponents δ , interpolating between the DP and original WPS exponents, $\delta = 0.4505(1)$ [2] and $\delta = 0.57(3)$ [39], respectively; Table I.

We performed extensive simulations of the CP on the modified WPS lattices with $L = \sqrt{N} = 8, 16, 32, ..., 512$,



FIG. 5. Density of active sites as a function of time in the decay analysis. The lines are power-law regressions (Pearson regression coefficient r > 0.999) of the data represented by symbols. An average over 100 lattice with 10 runs each was used.

where N = 3n + 1 is the number of nodes, employing the quasi-stationary (QS) simulation method [43]. It is based on maintaining and gradually updating a set of configurations visited during the evolution. If a transition to the absorbing state is imminent, the system is instead placed in one of the saved configurations. This procedure has been successfully applied to investigate critical dynamics on different disordered graphs [27, 28, 31, 40]. Each realization of the process was initialized with all sites occupied and run for at least 10⁶ time units. Averages were taken in the QS regime after discarding an initial transient. Typically, we performed averages over at least 100 distinct lattices for each size.



FIG. 6. Double-logarithm plots of the critical QS density of active sites ρ as a function of the $L = \sqrt{N}$ for different values of ϵ . Lines are linear regressions (Pearson regression coefficient r > 0.999) of data represented by symbols.



FIG. 7. Double-logarithm plots of the critical lifespan τ of the QS regime as a function of the $L = \sqrt{N}$ for different values of ϵ . Lines are linear regressions (Pearson regression coefficient r > 0.999) of data represented by symbols.

	λ_c	η	δ	eta/ u_{\perp}	z
DP	-	0.2295(10)	0.4505(1)	0.797(3)	1.7674(6)
$\epsilon = 0.10$	1.5703(2)	0.232(5)	0.45(1)	0.80(2)	1.76(3)
0.50	1.5675(2)	0.221(4)	0.47(2)	0.81(2)	1.68(4)
0.70	1.5637(1)	0.203(5)	0.50(1)	0.83(2)	1.60(3)
0.80	1.5611(1)	0.197(4)	0.52(2)	0.84(2)	1.58(3)
0.90	1.5577(1)	0.183(5)	0.54(2)	0.87(3)	1.56(3)
WPS	1.5525(5)	0.185(5)	0.57(1)	0.84(1)	1.59(1)

TABLE I. Critical rates and exponents for different values of the parameter ϵ . The uncertainties in last significant digit, indicated in parenthesis, were determined using a local slope analysis for the range $64 \leq L \leq 512$ for QS and $10^3 < t < 10^5$ for the dynamical quantities.

At $\lambda = \lambda_c$, the QS density of active sites, ρ , follows a power-law $\rho \sim L^{-\beta/\nu_{\perp}}$. Also, the lifetime of the QS state, τ , taken as the mean time between two attempts to visit the absorbing state in the QS simulation [43], follows $\tau \sim L^z$ at the critical point. The scaling laws presented in Figs. 6 and 7 lead to the exponents β/ν_{\perp} and z listed in Table I. Analogously to the time-dependent analysis, we observe that both exponents vary continuously with ϵ , from the values obtained for the DP class to the values for the original WPS lattice.

IV. CONCLUSIONS

We modified the algorithm proposed by Hassan et. al. [34] to construct a weighted planar stochastic lattice and investigate the role of planar disorder. The resulting lattice presents multifractality, and its dual network degree follows a power-law degree distribution $P(q) \sim q^{-\alpha}$ with continuously varying exponents interpolating between the WPS ($\alpha = 5.6$) and square lattice. We performed extensive large-scale simulations of the contact process on the modified WPS lattices and reported critical exponents depending on the parameter ϵ that controls the degree distribution of the graph. This result is qualitatively analogous to the contact processes on random, small-world, and thus infinite-dimensional graphs with power-law degree distribution [40, 41]. However, the relationship between the critical exponent and the degree distributions of the graphs is different. In the infinite-dimensional case, a mean-field approach where the degree heterogeneity is explicitly reckoned [40, 41] is sufficient to explain the critical exponent observed in simulations, while the CP in the planar, two-dimensional WPS networks presents the criticality at odds with the heterogeneous mean-field approach. In particular, the standard mean-field exponent, compatible with homogeneous networks, is recovered in the heterogeneous meanfield approach for $\alpha > 3$, whereas the simulation of CP on WPS lattices, which are featured by $\alpha > 5.6$, is far

- [1] J. Marro and R. Dickman, <u>Nonequilibrium Phase Transitions in Lattice Models</u> (Cambridge University Press, 1999).
- [2] M. Henkel, M. Pleimling, H. Hinrichsen, and S. Lübeck, <u>Non-Equilibrium Phase Transitions</u>, Vol. 2 (Springer Netherlands, 2008).
- [3] R. Pastor-Satorras, C. Castellano, P. V. Mieghem, and A. Vespignani, Epidemic processes in complex networks, Rev. Mod. Phys. 87, 925 (2015).
- [4] M. M. de Oliveira and R. Dickman, Quasi-stationary distributions for models of heterogeneous catalysis, Physica A 343, 10.1016/j.physa.2004.06.155 (2004).
- [5] R. M. Ziff, E. Gulari, and Y. Barshad, Kinetic phase transitions in an irreversible surface-reaction model, Phys. Rev. Lett. 56, 2553 (1986).
- [6] M. M. de Oliveira, R. V. Dos Santos, and R. Dickman, Symbiotic two-species contact process, Phys. Rev. E 86, 011121 (2012).
- [7] K. A. Takeuchi, M. Kuroda, H. Chaté, and M. Sano, Directed percolation criticality in turbulent liquid crystals, Phys. Rev. Lett. 99, 234503 (2007).
- [8] L. Corté, P. M. Chaikin, J. P. Gollub, and D. J. Pine, Random organization in periodically driven systems, Nat. Phys. 4, 420 (2008).
- [9] S. Okuma, Y. Tsugawa, and A. Motohashi, Transition from reversible to irreversible flow: Absorbing and depinning transitions in a sheared-vortex system, Phys. Rev. B 83, 012503 (2011).
- [10] H. K. Janssen, On the nonequilibrium phase transition in reaction-diffusion systems with an absorbing stationary state, Z. Phys. B 42, 151 (1981).
- [11] P. Grassberger, On phase transitions in schlögl's second model, Z. Phys. B 47, 365 (1982).
- [12] A. J. Noest, Power-law relaxation of spatially disordered stochastic cellular automata and directed percolation, Phys. Rev. B 38, 2715 (1988).
- [13] A. G. Moreira and R. Dickman, Critical dynamics of the contact process with quenched disorder, Phys. Rev. E 54,

from mean-field.

Finally, the analysis of degree fluctuations in the WPS lattices across different scales leads to an asymptotic wandering exponent $\omega = 1/2$ that indicates the disorder should be a relevant perturbation according to the Luck-Harris criterion [30]. Our work reveals agreement with the criterion, with weaker effects than usually observed for the DP class under uncorrelated disorder that presents the same wandering exponent for all strengths of the disorder controlled by the parameter $\epsilon \in [0.1, 0.9]$.

V. ACKNOWLEDGEMENT

SGA thanks the support of CNPq (Grant no. 311019/2021-8). MMO thanks the support of CNPq (Grant no. 312462/2021-2) and FAPEMIG (Grant no. APQ02393-18). SCF thanks the support of CNPq (Grant no. 310984/2023-8)

R3090 (1996).

- [14] R. Dickman and A. G. Moreira, Violation of scaling in the contact process with quenched disorder, Phys. Rev. E 57, 1263 (1998).
- [15] T. Vojta and M. Y. Lee, Nonequilibrium phase transition on a randomly diluted lattice, Phys. Rev. Lett. 96, 035701 (2006).
- [16] M. M. de Oliveira and S. C. Ferreira, Universality of the contact process with random dilution, J. Stat. Mech.: Theor. Exp., P11001 (2008).
- [17] T. Vojta, A. Farquhar, and J. Mast, Infinite-randomness critical point in the two-dimensional disordered contact process, Phys. Rev. E 79, 011111 (2009).
- [18] M. Bramson, R. Durrett, and R. H. Schonmann, The contact processes in a random environment, The Annals of Probability **19**, pp. 960 (1991).
- [19] M. S. Faria, D. J. Ribeiro, and S. R. Salinas, Critical behavior of a contact process with aperiodic transition rates, J. Stat. Mech.: Theor. Exp. **2008**, P01022 (2008).
- [20] T. Vojta and M. Dickison, Critical behavior and griffiths effects in the disordered contact process, Physical Review E 72, 036126 (2005).
- [21] T. Vojta, Rare region effects at classical, quantum and nonequilibrium phase transitions, Journal of Physics A: Mathematical and General **39**, R143 (2006).
- [22] M. Dickison and T. Vojta, Monte carlo simulations of the smeared phase transition in a contact process with extended defects, Journal of Physics A: Mathematical and General 38, 1199 (2005).
- [23] A. B. Harris, Effect of random defects on the critical behaviour of ising models, J. Phys. C 7, 1671 (1974).
- [24] J. Hooyberghs, F. Iglói, and C. Vanderzande, Strong disorder fixed point in absorbing-state phase transitions, Phys. Rev. Lett. **90**, 100601 (2003).
- [25] J. Hooyberghs, F. Iglói, and C. Vanderzande, Absorbing state phase transitions with quenched disorder, Phys. Rev. E 69, 066140 (2004).

- [26] A. Okabe, B. Boots, K. Sugihara, and S. N. Chiu, Spatial Tessellations: Concepts and Applications of Voronoi Diagdymamic scaling and neighbourhood statistics in edited by A. Okabe, B. Boots, K. Sugihara, S. N. Chiu, and D. G. Kendall (John Wiley & Sons, Inc., 2000).
- [27] M. M. de Oliveira, S. G. Alves, S. C. Ferreira, and R. Dickman, Contact process on a voronoi triangulation, Phys. Rev. E 78, 031133 (2008).
- [28] M. M. De Oliveira, S. Alves, and S. Ferreira, Continuous and discontinuous absorbing-state phase transitions on Voronoi-Delaunay random lattices, Physical Review E 93, 10.1103/PhysRevE.93.012110 (2016).
- [29] H. Barghathi and T. Vojta, Phase transitions on random lattices: How random is topological disorder?, Phys. Rev. Lett. 113, 120602 (2014).
- [30] J. M. Luck, A classification of critical phenomena on quasi-crystals and other aperiodic structures, Europhysics Letters (EPL) 24, 359 (1993).
- [31] M. Schrauth, J. S. Portela, and F. Goth, Violation of the Harris-Barghathi-Vojta Criterion, Physical Review Letters 121, 100601 (2018), 1803.08753.
- [32] M. Schrauth and J. S. E. Portela, Universality of continuous phase transitions on random voronoi graphs, Phys. Rev. E **100**, 062118 (2019).
- [33] M. Schrauth and J. S. E. Portela, Fast algorithm for topologically disordered lattices with constant coordination number, Phys. Rev. Research 1, 033061 (2019).
- [34] M. K. Hassan, M. Z. Hassan, and N. I. Pavel, Scale-free network topology and multifractality in a weighted planar stochastic lattice, New Journal of Physics 12, 093045 (2010).

- [35] F. R. Dayeen and M. K. Hassan, Multi-multifractality,
- weighted planar stochastic lattice, Chaos, Solitons and Fractals 91, 228 (2016).
- [36] M. K. Hassan and M. M. Rahman, Universality class of site and bond percolation on multifractal scale-free planar stochastic lattice, Phys. Rev. E 94, 042109 (2016).
- [37] H.-P. Hsu and M.-C. Huang, Percolation thresholds, critical exponents, and scaling functions on planar random lattices and their duals, Phys. Rev. E 60, 6361 (1999).
- X.-S. Liu, Z.-X. Wu, and J.-Y. Guan, Kinetic-exchange-[38]like opinion dynamics in complex networks: roles of the dimensionality and local interaction topology, The European Physical Journal B **91**, 220 (2018).
- [39] S. G. Alves and M. M. de Oliveira, Contact process on weighted planar stochastic lattice, J. Stat. Mech.: Theor. Exp. 2022, 063201 (2022).
- [40] S. C. Ferreira, R. S. Ferreira, C. Castellano, and R. Pastor-Satorras, Quasistationary simulations of the contact process on quenched networks, Phys. Rev. E 84, 066102 (2011).
- [41] A. S. Mata, R. S. Ferreira, and S. C. Ferreira, Heterogeneous pair-approximation for the contact process on complex networks, New J. Phys. 16, 053006 (2014).
- [42]W. Cota and S. C. Ferreira, Optimized gillespie algorithms for the simulation of markovian epidemic processes on large and heterogeneous networks, Computer Physics Communications 219, 303 (2017).
- [43] M. M. de Oliveira and R. Dickman, How to simulate the quasistationary state, Phys. Rev. E 71, 016129 (2005).